

United States Patent Application

of

Dr. Uwe Heinelt  
Dr. Hans-Jochen Lang  
Dr. Klaus Wirth  
and  
Dr. Hans-Willi Jansen

for

**Substituted heterocyclo-norbornylamino derivatives, processes for  
their preparation, their use as medicaments or diagnostics, and  
medicaments comprising them**


FILED

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[001] This application claims the benefit of the filing date of German Patent Application Number 10063294.7-44, filed on December 19, 2000, which application is hereby incorporated by reference.

[002] One embodiment of the invention relates to substituted heterocyclo-norbornylamino derivatives having: a) an exo-configuration nitrogen and an endo-fused five-membered or six-membered ring of the formula (I), or b) an exo-configuration nitrogen and an exo-fused five-membered or six-membered ring of the formula (I a),



wherein:

[003] A is (C<sub>1</sub>-C<sub>4</sub>)-alkylene;

[004] T is (C<sub>1</sub>-C<sub>4</sub>)-alkyl or H;

[005] B is a saturated or unsaturated carbon five-membered or six-membered ring, which is unsubstituted or is substituted, having 1-3 substituents chosen from oxo, hydroxyl, (C<sub>1</sub>-C<sub>4</sub>)-alkoxy, and (C<sub>1</sub>-C<sub>4</sub>)-alkyl;

[006] Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to four identical or different heteroatoms chosen from O, S, N, and Se;

[007] R is OH, F, Cl, Br, I, CN, NO<sub>2</sub>, phenyl, CO<sub>2</sub>R<sub>1</sub>, (C<sub>1</sub>-C<sub>4</sub>)-alkyl, (C<sub>1</sub>-C<sub>4</sub>)-alkoxy, amino, (C<sub>1</sub>-C<sub>4</sub>)-alkylamino, di(C<sub>1</sub>-C<sub>4</sub>)-alkylamino, or amino-(C<sub>1</sub>-C<sub>4</sub>)-alkyl,

[008] wherein the alkyl radicals are unsubstituted or are completely or partly substituted by fluorine;

[009] R<sub>1</sub> is H or (C<sub>1</sub>-C<sub>4</sub>)-alkyl, which is unsubstituted or completely or partly substituted by fluorine;

[010] n0, 1, 2, 3 or 4,

[011] wherein, if n = 2, 3 or 4, the substituents R are chosen independently of one another;

[012] and their pharmaceutically tolerable salts or trifluoracetates.

[013] Examples of compounds of the invention include those compounds having an exo-configuration nitrogen and endo-fused carbon five-membered or six-membered ring of the formula (I) and those compounds having an exo-configuration nitrogen and exo-fused carbon five-membered or six-membered ring of the formula (I a), wherein:

[014] A is (C<sub>1</sub>-C<sub>2</sub>)- alkylene;

[015] T is H or methyl;

[016] B is a saturated or unsaturated carbon five-membered or six-membered ring;

[017] Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to three identical or different heteroatoms chosen from O, S, and N;

[018] R is F, Cl, Br, iodine, amino, hydroxymethyl, OH, phenyl, CO<sub>2</sub>R<sub>1</sub>, (C<sub>1</sub>-C<sub>4</sub>)- alkyl, or (C<sub>1</sub>-C<sub>4</sub>)- alkoxy,

[019] wherein the alkyl radicals are unsubstituted or completely or partly substituted by fluorine;

[020] R<sub>1</sub> is H or (C<sub>1</sub>-C<sub>4</sub>)-alkyl, wherein the alkyl radical is unsubstituted or completely or partly substituted by fluorine;

[021] n0, 1, 2 or 3,

[022] wherein, if n = 2 or 3, the corresponding substituents R are chosen independently of one another;

[023] and their pharmaceutically tolerable salts or trifluoracetates.

[024] Further examples of compounds of the invention include those compounds having an exo-configuration nitrogen and endo-fused carbon five-membered or six-membered ring of the formula (I) and those compounds having an

exo-configuration nitrogen and exo-fused carbon five-membered or six-membered ring of the formula (I a), wherein:

[025] A is (C<sub>1</sub>-C<sub>2</sub>)- alkylene;

[026] T is hydrogen;

[027] B is a saturated or unsaturated carbon five-membered or six-membered ring;

[028] Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to two identical or different heteroatoms chosen from O, S, and N ;

[029] R is F, Cl, Br, (C<sub>1</sub>-C<sub>4</sub>)- alkoxy or (C<sub>1</sub>-C<sub>4</sub>)- alkyl,

[030] wherein the alkyl radicals are unsubstituted or completely or partly substituted by fluorine;

[031] n is 0, 1 or 2, wherein, if n = 2, the corresponding substituents R are chosen independently of one another;

[032] and their pharmaceutically tolerable salts or trifluoroacetates.

[033] Even more examples of the compounds of the invention include those compounds having an exo-configuration nitrogen and endo-fused carbon five-membered or six-membered ring of the formula (I) and those compounds having an exo-configuration nitrogen and exo-fused carbon five-membered ring of the formula (I a) such as, for example:

exo/exo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,  
 (rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,  
 (+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,  
 (-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,  
 (rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,  
 (+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,  
 (-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,  
 exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)pyrazin-2-ylmethylamine,  
 exo/endo-(octahydro-4,7-methanoinden-5-yl)thiophen-2-ylmethylamine,  
 exo/endo-(octahydro-4,7-methanoinden-5-yl)thiophen-3-ylmethylamine,

[illegible]

**LAW OFFICES**  
**EGAN, HENDERSON,**  
**LABOW, GARRETT,**  
**DUNNER, L.L.P.**  
**101 STREET, N. W.**  
**WASHINGTON, DC 20005**  
**202-408-4000**

- 4 -


and their pharmaceutically tolerable salts or trifluoroacetates.

[035] Suitable acid addition salts of the compounds of the invention include salts of pharmacologically tolerable acids, for example halides (such as hydrochlorides), lactates, sulfates, citrates, tartrates, acetates, phosphates, methylsulfonates, p-toluenesulfonates, adipates, fumarates, gluconates, glutamates, glycerophosphates, maleates, and pamoates. This group also exemplifies some of the physiologically acceptable anions, in addition to trifluoroacetates.

[036] If a compound of the formula (I) or (I a) contains one or more asymmetric centers, these centers can have either the S or the R configuration. The compounds can be present as optical isomers, diastereomers, racemates, or mixtures thereof. However, the amino substituent on the norbornyl system must be exo and the ring endo- or exo-fused



[037] The alkyl or alkylene radicals mentioned can be either straight-chain or branched.

[038] Suitable heterocycles include, *inter alia*:




[039] Another embodiment of the invention relates to a process for the preparation of the compounds of the formula (I) or (I a), which comprises

[040] a) reacting a compound of the formula (II) or (II a)





[041] with a compound of the formula (III)



[042] in the presence of suitable reductants and possibly also Lewis acids directly to give compounds of the formula (I) or (I a),

[043] wherein T, B, Het and R<sub>n</sub> have the meanings indicated above; independently of one another A' corresponds to a bond or (C<sub>1</sub>-C<sub>3</sub>)- alkyl and A'' corresponds to H or (C<sub>1</sub>-C<sub>3</sub>)- alkyl; and A' and A'', together with the carbon atom of the carbonyl group, represent as many carbon atoms as A represents in formula (I) or (I a); or

[044] b) isolating an intermediate of the formula (IV) or (IV a), formed from reacting compounds of the formulae (II), or (II a), with a compound of formula (III), and then converting the intermediate of the formula (IV) or (IV a) into the compounds of the formula (I) or (I a) by using suitable reductants,




LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[045] wherein T# is a free electron pair or (C<sub>1</sub>-C<sub>4</sub>)-alkyl. When T# is (C<sub>1</sub>-C<sub>4</sub>)-alkyl, an iminium ion is formed, to which a counterion is assigned, such as, for example, chloride or tosylate,

or



[046] c) reacting a compound of the formula (II) or (II a) with an alkylating agent of the formula (V),



[047] wherein U is a nucleophilically substitutable group, such as for example a halogen, alkylsulfonates, or arylsulfonates, including Cl, Br, I, mesylate, or tosylate, and the other radicals are defined as described above. Here, the carbon atom to which U is bonded corresponds to the carbon atom of the carbonyl group of the compound of formula (III),

or

[048] d) reducing carboxamides of the formula (VI) or (VI a) to the corresponding amines,



[049] wherein A\* corresponds to a bond or (C<sub>1</sub>-C<sub>3</sub>)-alkyl and the other radicals are defined as described above,

or




[050] e) alkylating compounds of the formula (I) or (I a) , in which T corresponds to hydrogen, using alkylating agents of the formula (VII),




[051] wherein T\* is(C<sub>1</sub>-C<sub>4</sub>)- alkyl and U has the meaning described above, so that tertiary amines result from this reaction;

or

[052] f) reacting a dicyclopentadienylplatinum complex of the formula (VIII)



[053] with amines of the formula (IX),



[054] and subsequently reducing the intermediate formed to compounds of the formula (I) or (I a) (J. K. Stille and D. B. Fox, JACS 92:1274 (1970)),

[055] wherein T, R<sub>n</sub> and Het have the meanings indicated above; independently of one another, A' corresponds to a bond or (C<sub>1</sub>-C<sub>3</sub>)-alkyl and A'' to H or (C<sub>1</sub>-C<sub>3</sub>)-alkyl; and A' and A'', together with the carbon atom to which the nitrogen atom is bonded, represent as many carbon atoms as A represents in formula (I).

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[056] The compounds of formula (I) or (I a) may be optionally converted into the pharmaceutically tolerable salt or trifluoroacetate.

[057] It has already been proposed that phenylalkyl-substituted norbornylamino derivatives are effective inhibitors of the sodium-proton exchanger, subtype 3 (NHE3). In this case, it appears that, of several stereoisomers, the compounds having an exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine unit, wherein the nitrogen is exo and the five-membered ring is endo-fused, are active NHE3 inhibitors. Substances having an exo/exo-configuration octahydro-4,7-methanoinden-5-ylamine unit likewise showed marked NHE3-inhibiting action, while the corresponding endo/endo and endo/exo-derivatives were markedly less active on the NHE3 (German published application 199 60 204 A1 - HMR 99/L 073).

[058] Surprisingly, it has now been found that the aromatic moiety of the phenylalkyl substituents can be substituted by heteroaromatic rings producing NHE3-inhibiting activity.

[059] The relatively long-known inhibitors of the sodium/proton exchanger, subtype 3 disclosed in EP-A 825 178 (HOE 96/F226) represent relatively polar structures and correspond to the acylguanidine type of compounds (J.-R. Schwark et al. Eur. J. Physiol (1998) 436:797). In contrast, the compounds according to the invention are surprisingly lipophilic substances that are not of the acylguanidine type. See also the proposed compounds of the phenylalkyl norbornylamine type (DE 199 60 204.2 - HMR 99 / L 073). Squalamine and the above phenylalkyl norbornylamino derivatives are only the fourth substance class of NHE3 inhibitors known hitherto (M. Donowitz *et al.*, Am. J. Physiol. 276 (Cell Physiol. 45): C136-C144). Additionally, squalamine does not achieve its maximum potency immediately, but only after approximately one hour. Compared with the above acylguanidines, the compounds of the invention are distinguished by their superior ability to cross the membrane and, compared with squalamine, by a more rapid onset of action.

0  
1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2

[060] NHE3 is found in the body of various species, mostly in the bile, the intestine, and in the kidney (Larry Fliegel *et al.*, Biochem. Cell. Biol. 76: 735-741 (1998)), but can also be detected in the brain (E. Ma *et al.*, Neuroscience 79: 591-603).

[061] The compounds of the formula (I) or (I a) according to the invention are suitable for use as antihypertensives for the treatment of primary and secondary hypertension.

[062] Moreover, these compounds, on their own or in combination with NHE inhibitors of other subtype specificity, can protect acutely or chronically oxygen-deficiently-supplied organs by reducing or preventing ischemically induced damage. Thus, the compounds of the invention are suitable as pharmaceuticals, in the treatment of acute or chronic kidney failure and during surgical interventions such as in organ transplantation of the kidney and liver, where the compounds can be used both for the protection of the organs in the donor before and during removal and for the protection of removed organs, (for example during treatment with or storage in physiological bath fluids, and during transfer to the recipient's body). Ischemically induced damage to the intestine can also be avoided.

[063] Corresponding to the protective action against ischemically induced damage, the compounds of the invention are also potentially suitable as pharmaceuticals for the treatment of ischemia of the nervous system, including the CNS, where they are suitable, for example, for the treatment of stroke or cerebral odema. Moreover, the compounds of the formula (I) or (I a) according to the invention are likewise suitable for the treatment of forms of shock, such as, for example, allergenic, cardiogenic, hypovolemic, and bacterial shock.

[064] Furthermore, the compounds of the invention induce an improvement in the respiratory drive and are therefore used for the treatment of respiratory conditions in the following clinical conditions and diseases: disturbed central respiratory drive (e.g.

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

central sleep apnea, sudden infant death, post operative hypoxia), muscular-related respiratory disorders, respiratory disorders after long-term ventilation, respiratory disorders during adaptation in a high mountain area, obstructive and mixed forms of sleep apnea, acute and chronic lung diseases with hypoxia and hypocapnia.

[065] The compounds of the invention additionally increase the muscle tone of the upper airways, so that snoring is suppressed.

[066] A combination of an NHE inhibitor with a carboanhydrase inhibitor, (e. g. acetazolamide) where the latter produces metabolic acidosis and thereby even increases the respiratory activity, proves to be advantageous as a result of increased activity and decreased use of active compound.

[067] It has been shown that the compounds according to the invention have a mild laxative action and accordingly can be used advantageously as laxatives or, in the case of threatening intestinal blockage, for the prevention of ischemic damage that accompanies blockages in the intestinal region.

[068] Furthermore, the use of the compounds of the formula (I) or (I a) according to the invention makes it possible to prevent gallstone formation.

[069] The compounds of the formula (I) or (I a) according to the invention additionally show an action against ectoparasites.

[070] Moreover, the compounds of the formula (I) or (I a) according to the invention can exert an inhibitory action on the proliferation of cells, for example fibroblast cell proliferation and the proliferation of the smooth vascular muscle cells. The compounds of the formula (I) or (I a) are therefore suitable as valuable therapeutics for diseases in which cell proliferation is a primary or secondary cause, and can therefore be used as antiatherosclerotics, as agents against diabetic late complications, cancers,

2481.1763

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

fibrotic disorders (such as, for example, pulmonary fibrosis, liver fibrosis, or kidney fibrosis), endothelial dysfunction, and organ hypotrophy and hyperplasia, such as, for example, prostate hyperplasia or prostate hypertrophy.

[071] The compounds according to the invention are effective inhibitors of the cellular sodium/proton antiporter, which is raised in numerous disorders (essential hypertension, atherosclerosis, diabetes etc.) even in those cells that are easily accessible to measurement, such as, for example, erythrocytes, platelets or leukocytes. The compounds according to the invention are therefore suitable as outstanding and simple scientific tools, for example in their use as diagnostics for the determination and differentiation of certain forms of hypertension, atherosclerosis, diabetes, proliferative disorders, etc. Moreover, the compounds of the formula (I) or (I a) are suitable for preventive therapy to prevent the development of high blood pressure, for example of essential hypertension.

[072] It has moreover been found that NHE inhibitors exhibit a favorable influence on the serum lipoproteins. It is generally recognized that for the formation of arteriosclerotic vascular changes, for example in coronary heart disease, excessively high blood lipid values, 'hyperlipoproteinemias', are an important risk factor. The lowering of increased serum lipoproteins is therefore of extreme importance for the prophylaxis and the regression of atherosclerotic changes. The compounds according to the invention can therefore be used for the prophylaxis and regression of atherosclerotic changes by excluding a causal risk factor. With this protection of the vessels against the syndrome of endothelial dysfunction, compounds of the formula (I) or (I a) are valuable pharmaceuticals for the prevention and treatment of coronary vascular spasms, atherogenesis and atherosclerosis, left-ventricular hypertrophy and dilated cardiomyopathy, and thrombotic disorders.

[073] The compounds mentioned are therefore advantageously used for the production of a medicament for the prevention and treatment of sleep apneas and

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N.W.  
WASHINGTON, DC 20005  
202-408-4000

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N.W.  
WASHINGTON, DC 20005  
202-408-4000

muscular-related respiratory disorders; for the production of a medicament for the prevention and treatment of snoring, for the production of a medicament for lowering the blood pressure, for the production of a medicament having laxative action for the prevention and treatment of intestinal blockages; for the production of a medicament for the prevention and treatment of disorders that are induced by ischemia and reperfusion of central and peripheral organs, such as acute kidney failure, stroke, endogenous states of shock, intestinal disorders, etc; for the production of a medicament for the treatment of hypercholesterolemia; for the production of a medicament for the prevention of atherogenesis and of athereosclerosis; for the production of a medicament for the prevention and treatment of diseases which are induced by raised cholesterol levels; for the production of a medicament for the prevention and treatment of diseases which are induced by endothelial dysfunction; for the production of a medicament for the treatment of attack by ectoparasites; for the production of a medicament for the treatment of the diseases mentioned in combination with blood pressure-lowering substances, for example angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor antagonists. A combination of an NHE inhibitor of the formula (I) or (I a) with an active compound lowering the blood lipid level, such as an HMG-CoA reductase inhibitor (e.g. Lovastatin or Pravastatin) proves to be a favorable combination having intensified action and decreased use of active substance. In this case, the HMG-CoA reductase inhibitor produces a hypolipodemic effect and thereby increases the hypolipodemic properties of the NHE inhibitor of the formula (I) or (I a).

[074] The administration of sodium/proton exchange inhibitors of the formula (I) or (I a) as novel pharmaceuticals for lowering raised blood lipid levels, and the combination of sodium/proton exchange inhibitors with pharmaceuticals having a blood pressure-lowering and/or hypolipodemic action is claimed.

[075] Pharmaceuticals containing a compound of formula (I) or (I a) can be administered orally, parenterally, intravenously, rectally, or by inhalation. The method of administration depends on the particular clinical picture of the disorder. The compounds

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

(I) or (I a) can be used on their own or together with pharmaceutical excipients, namely both in veterinary and in human medicine.

[076] The person skilled in the art is familiar on the basis of his/her expert knowledge with excipients that are suitable for the desired pharmaceutical formulation. In addition to solvents, gel formers, suppository bases, tablet excipients, and other active compound carriers, it is possible to use, for example, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, and/or colorants.

[077] For a form suitable for oral administration, the active compounds are ~~mixed with the additives suitable~~ therefor, including excipients, stabilizers, or inert diluents, and brought by means of the customary methods into the suitable administration forms, including tablets, coated tablets, hard gelatin capsules, aqueous, alcoholic, or oily solutions. Inert carriers that can be used include, for example, gum arabic, magnesia, magnesium carbonates, potassium phosphates, lactose, glucose, starch, or cornstarch. Preparation can be carried out either as dry or as moist granules. Possible oily excipients or solvents include, for example, vegetable or animal oils, such as, for example, sunflower oil or cod-liver oils.

[078] For subcutaneous or intravenous administration, the active compounds are brought into solution, suspension, or emulsion, if desired with the substances customary therefor such as, for example, solubilizers, emulsifiers, or further excipients. Suitable solvents include, for example, water, physiological saline solution, or alcohols (e.g. ethanol, propanol, glycerol), and sugar solutions such, for example, as glucose or mannitol solutions, or alternatively a mixture of the different solvents mentioned.

[079] Pharmaceutical formulations suitable for administration in the form of aerosols or sprays include, for example, solutions, suspensions, or emulsions of the active compound of the formula (I) or (I a) in a pharmaceutically innocuous solvent, such as, ethanol or water, or a mixture of such solvents.

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[080] If required, the formulation can also contain other pharmaceutical excipients such as, for example, surfactants, emulsifiers and stabilizers, and a propellant. Such a preparation may contain the active compound in a concentration of approximately 0.1 to 10 % by weight; and sometimes approximately 0.3 to 3 % by weight.

[081] The dose of the active compound of the formula (I) or (I a) to be administered and the frequency of administration depend on the potency and duration of action of the compounds used; on the nature and severity of the disease to be treated; and on the sex, age, weight, and individual responsiveness of the mammal to be treated.

[082] On average, the daily dose of a compound of the formula (I) or (I a) in the case of a patient weighing approximately 75 kg may be at least 0.001 mg/kg, sometimes 1 - 10 mg/kg, and at most 100 mg/kg, of bodyweight. In acute episodes of the diseases, even higher and more frequent doses may also be necessary, e.g. up to 4 individual doses per day, for instance in the case of i.v. administration, for example in the case of an infarct patient in the intensive care unit, up to 200 mg per day may be necessary.

#### Experimental section:

[083] Abbreviations used:

CI	chemical ionization
DIP	diisopropyl ether
EA	ethyl acetate
ES	electrospray
FAB	fast atom bombardment
h	hours

bp	boiling point
MgSO <sub>4</sub>	magnesium sulfate
m.p.	melting point
MS	mass spectrum
NaOH	sodium hydroxide solution
RP	reversed phase

11-12-13-14-15-16-17-18-19-20-21-22-23-24-25-26-27-28-29-30-31-32-33-34-35-36-37-38-39-40-41-42-43-44-45-46-47-48-49-50-51-52-53-54-55-56-57-58-59-60-61-62-63-64-65-66-67-68-69-70-71-72-73-74-75-76-77-78-79-80-81-82-83-84-85-86-87-88-89-90-91-92-93-94-95-96-97-98-99-100-101-102-103-104-105-106-107-108-109-110-111-112-113-114-115-116-117-118-119-120-121-122-123-124-125-126-127-128-129-130-131-132-133-134-135-136-137-138-139-140-141-142-143-144-145-146-147-148-149-150-151-152-153-154-155-156-157-158-159-160-161-162-163-164-165-166-167-168-169-170-171-172-173-174-175-176-177-178-179-180-181-182-183-184-185-186-187-188-189-190-191-192-193-194-195-196-197-198-199-200-201-202-203-204-205-206-207-208-209-210-211-212-213-214-215-216-217-218-219-220-221-222-223-224-225-226-227-228-229-230-231-232-233-234-235-236-237-238-239-240-241-242-243-244-245-246-247-248-249-250-251-252-253-254-255-256-257-258-259-260-261-262-263-264-265-266-267-268-269-270-271-272-273-274-275-276-277-278-279-280-281-282-283-284-285-286-287-288-289-290-291-292-293-294-295-296-297-298-299-300-301-302-303-304-305-306-307-308-309-310-311-312-313-314-315-316-317-318-319-320-321-322-323-324-325-326-327-328-329-330-331-332-333-334-335-336-337-338-339-340-341-342-343-344-345-346-347-348-349-350-351-352-353-354-355-356-357-358-359-360-361-362-363-364-365-366-367-368-369-370-371-372-373-374-375-376-377-378-379-380-381-382-383-384-385-386-387-388-389-390-391-392-393-394-395-396-397-398-399-400-401-402-403-404-405-406-407-408-409-410-411-412-413-414-415-416-417-418-419-420-421-422-423-424-425-426-427-428-429-430-431-432-433-434-435-436-437-438-439-440-441-442-443-444-445-446-447-448-449-450-451-452-453-454-455-456-457-458-459-460-461-462-463-464-465-466-467-468-469-470-471-472-473-474-475-476-477-478-479-480-481-482-483-484-485-486-487-488-489-490-491-492-493-494-495-496-497-498-499-500-501-502-503-504-505-506-507-508-509-510-511-512-513-514-515-516-517-518-519-520-521-522-523-524-525-526-527-528-529-530-531-532-533-534-535-536-537-538-539-540-541-542-543-544-545-546-547-548-549-550-551-552-553-554-555-556-557-558-559-560-561-562-563-564-565-566-567-568-569-570-571-572-573-574-575-576-577-578-579-580-581-582-583-584-585-586-587-588-589-590-591-592-593-594-595-596-597-598-599-600-601-602-603-604-605-606-607-608-609-610-611-612-613-614-615-616-617-618-619-620-621-622-623-624-625-626-627-628-629-630-631-632-633-634-635-636-637-638-639-640-641-642-643-644-645-646-647-648-649-650-651-652-653-654-655-656-657-658-659-660-661-662-663-664-665-666-667-668-669-670-671-672-673-674-675-676-677-678-679-680-681-682-683-684-685-686-687-688-689-690-691-692-693-694-695-696-697-698-699-700-701-702-703-704-705-706-707-708-709-710-711-712-713-714-715-716-717-718-719-720-721-722-723-724-725-726-727-728-729-730-731-732-733-734-735-736-737-738-739-740-741-742-743-744-745-746-747-748-749-750-751-752-753-754-755-756-757-758-759-760-761-762-763-764-765-766-767-768-769-770-771-772-773-774-775-776-777-778-779-780-781-782-783-784-785-786-787-788-789-790-791-792-793-794-795-796-797-798-799-800-801-802-803-804-805-806-807-808-809-810-811-812-813-814-815-816-817-818-819-820-821-822-823-824-825-826-827-828-829-830-831-832-833-834-835-836-837-838-839-840-841-842-843-844-845-846-847-848-849-850-851-852-853-854-855-856-857-858-859-860-861-862-863-864-865-866-867-868-869-870-871-872-873-874-875-876-877-878-879-880-881-882-883-884-885-886-887-888-889-890-891-892-893-894-895-896-897-898-899-900-901-902-903-904-905-906-907-908-909-910-911-912-913-914-915-916-917-918-919-920-921-922-923-924-925-926-927-928-929-930-931-932-933-934-935-936-937-938-939-940-941-942-943-944-945-946-947-948-949-950-951-952-953-954-955-956-957-958-959-960-961-962-963-964-965-966-967-968-969-970-971-972-973-974-975-976-977-978-979-980-981-982-983-984-985-986-987-988-989-990-991-992-993-994-995-996-997-998-999-1000-1001-1002-1003-1004-1005-1006-1007-1008-1009-1010-1011-1012-1013-1014-1015-1016-1017-1018-1019-1020-1021-1022-1023-1024-1025-1026-1027-1028-1029-1030-1031-1032-1033-1034-1035-1036-1037-1038-1039-1040-1041-1042-1043-1044-1045-1046-1047-1048-1049-1050-1051-1052-1053-1054-1055-1056-1057-1058-1059-1060-1061-1062-1063-1064-1065-1066-1067-1068-1069-1070-1071-1072-1073-1074-1075-1076-1077-1078-1079-1080-1081-1082-1083-1084-1085-1086-1087-1088-1089-1090-1091-1092-1093-1094-1095-1096-1097-1098-1099-1100-1101-1102-1103-1104-1105-1106-1107-1108-1109-1110-1111-1112-1113-1114-1115-1116-1117-1118-1119-1120-1121-1122-1123-1124-1125-1126-1127-1128-1129-1130-1131-1132-1133-1134-1135-1136-1137-1138-1139-1140-1141-1142-1143-1144-1145-1146-1147-1148-1149-1150-1151-1152-1153-1154-1155-1156-1157-1158-1159-1160-1161-1162-1163-1164-1165-1166-1167-1168-1169-1170-1171-1172-1173-1174-1175-1176-1177-1178-1179-1180-1181-1182-1183-1184-1185-1186-1187-1188-1189-1190-1191-1192-1193-1194-1195-1196-1197-1198-1199-1200-1201-1202-1203-1204-1205-1206-1207-1208-1209-1210-1211-1212-1213-1214-1215-1216-1217-1218-1219-1220-1221-1222-1223-1224-1225-1226-1227-1228-1229-1230-1231-1232-1233-1234-1235-1236-1237-1238-1239-1240-1241-1242-1243-1244-1245-1246-1247-1248-1249-1250-1251-1252-1253-1254-1255-1256-1257-1258-1259-1260-1261-1262-1263-1264-1265-1266-1267-1268-1269-1270-1271-1272-1273-1274-1275-1276-1277-1278-1279-1280-1281-1282-1283-1284-1285-1286-1287-1288-1289-1290-1291-1292-1293-1294-1295-1296-1297-1298-1299-1300-1301-1302-1303-1304-1305-1306-1307-1308-1309-1310-1311-1312-1313-1314-1315-1316-1317-1318-1319-1320-1321-1322-1323-1324-1325-1326-1327-1328-1329-1330-1331-1332-1333-1334-1335-1336-1337-1338-1339-1340-1341-1342-1343-1344-1345-1346-1347-1348-1349-1350-1351-1352-1353-1354-1355-1356-1357-1358-1359-1360-1361-1362-1363-1364-1365-1366-1367-1368-1369-1370-1371-1372-1373-1374-1375-1376-1377-1378-1379-1380-1381-1382-1383-1384-1385-1386-1387-1388-1389-1390-1391-1392-1393-1394-1395-1396-1397-1398-1399-1400-1401-1402-1403-1404-1405-1406-1407-1408-1409-1410-1411-1412-1413-1414-1415-1416-1417-1418-1419-1420-1421-1422-1423-1424-1425-1426-1427-1428-1429-1430-1431-1432-1433-1434-1435-1436-1437-1438-1439-1440-1441-1442-1443-1444-1445-1446-1447-1448-1449-1450-1451-1452-1453-1454-1455-1456-1457-1458-1459-1460-1461-1462-1463-1464-1465-1466-1467-1468-1469-1470-1471-1472-1473-1474-1475-1476-1477-1478-1479-1480-1481-1482-1483-1484-1485-1486-1487-1488-1489-1490-1491-1492-1493-1494-1495-1496-1497-1498-1499-1500-1501-1502-1503-1504-1505-1506-1507-1508-1509-1510-1511-1512-1513-1514-1515-1516-1517-1518-1519-1520-1521-1522-1523-1524-1525-1526-1527-1528-1529-1530-1531-1532-1533-1534-1535-1536-1537-1538-1539-1540-1541-1542-1543-1544-1545-1546-1547-1548-1549-1550-1551-1552-1553-1554-1555-1556-1557-1558-1559-1560-1561-1562-1563-1564-1565-1566-1567-1568-1569-1570-1571-1572-1573-1574-1575-1576-1577-1578-1579-1580-1581-1582-1583-1584-1585-1586-1587-1588-1589-1590-1591-1592-1593-1594-1595-1596-1597-1598-1599-1600-1601-1602-1603-1604-1605-1606-1607-1608-1609-1610-1611-1612-1613-1614-1615-1616-1617-1618-1619-1620-1621-1622-1623-1624-1625-1626-1627-1628-1629-1630-1631-1632-1633-1634-1635-1636-1637-1638-1639-1640-1641-1642-1643-1644-1645-1646-1647-1648-1649-1650-1651-1652-1653-1654-1655-1656-1657-1658-1659-1660-1661-1662-1663-1664-1665-1666-1667-1668-1669-1670-1671-1672-1673-1674-1675-1676-1677-1678-1679-1680-1681-1682-1683-1684-1685-1686-1687-1688-1689-1690-1691-1692-1693-1694-1695-1696-1697-1698-1699-1700-1701-1702-1703-1704-1705-1706-1707-1708-1709-1710-1711-1712-1713-1714-1715-1716-1717-1718-1719-1720-1721-1722-1723-1724-1725-1726-1727-1728-1729-1730-1731-1732-1733-1734-1735-1736-1737-1738-1739-1740-1741-1742-1743-1744-1745-1746-1747-1748-1749-1750-1751-1752-1753-1754-1755-1756-1757-1758-1759-1760-1761-1762-1763-1764-1765-1766-1767-1768-1769-1770-1771-1772-1773-1774-1775-1776-1777-1778-1779-1780-1781-1782-1783-1784-1785-1786-1787-1788-1789-1790-1791-1792-1793-1794-1795-1796-1797-1798-1799-1800-1801-1802-1803-1804-1805-1806-1807-1808-1809-1810-1811-1812-1813-1814-1815-1816-1817-1818-1819-1820-1821-1822-1823-1824-1825-1826-1827-1828-1829-1830-1831-1832-1833-1834-1835-1836-1837-1838-1839-1840-1841-1842-1843-1844-1845-1846-1847-1848-1849-1850-1851-1852-1853-1854-1855-1856-1857-1858-1859-1860-1861-1862-1863-1864-1865-1866-1867-1868-1869-1870-1871-1872-1873-1874-1875-1876-1877-1878-1879-1880-1881-1882-1883-1884-1885-1886-1887-1888-1889-1890-1891-1892-1893-1894-1895-1896-1897-1898-1899-1900-1901-1902-1903-1904-1905-1906-1907-1908-1909-1910-1911-1912-1913-1914-1915-1916-1917-1918-1919-1920-1921-1922-1923-1924-1925-1926-1927-1928-1929-1930-1931-1932-1933-1934-1935-1936-1937-1938-1939-1940-1941-1942-1943-1944-1945-1946-1947-1948-1949-1950-1951-1952-1953-1954-1955-1956-1957-1958-1959-1960-1961-1962-1963-1964-1965-1966-1967-1968-1969-1970-1971-1972-1973-1974-1975-1976-1977-1978-1979-1980-1981-1982-1983-1984-1985-1986-1987-1988-1989-1990-1991-1992-1993-1994-1995-1996-1997-1998-1999-2000-2001-2002-2003-2004-2005-2006-2007-2008-2009-2010-2011-2012-2013-2014-2015-2016-2017-2018-2019-2020-2021-2022-2023-2024-2025-2026-2027-2028-2029-2030-2031-2032-2033-2034-2035-2036-2037-2038-2039-2040-2041-2042-2043-2044-2045-2046-2047-2048-2049-2050-2051-2052-2053-2054-2055-2056-2057-2058-2059-2060-2061-2062-2063-2064-2065-2066-2067-2068-2069-2070-2071-2072-2073-2074-2075-2076-2077-2078-2079-2080-2081-2082-2083-2084-2085-2086-2087-2088-2089-2090-2091-2092-2093-2094-2095-2096-2097-2098-2099-2100-2101-2102-2103-2104-2105-2106-2107-2108-2109-2110-2111-2112-2113-2114-2115-2116-2117-2118-2119-2120-2121-2122-2123-2124-2125-2126-2127-2128-2129-2130-2131-2132-2133-2134-2135-2136-2137-2138-2139-2140-2141-2142-2143-2144-2145-2146-2147-2148-2149-2150-2151-2152-2153-2154-2155-2156-2157-2158-2159-2160-2161-2162-2163-2164-2165-2166-2167-2168-2169-2170-2171-2172-2173-2174-2175-2176-2177-2178-2179-2180-2181-2182-2183-2184-2185-2186-2187-2188-2189-2190-2191-2192-2193-2194-2195-2196-2197-2198-2199-2200-2201-2202-2203-2204-2205-2206-2207-2208-2209-2210-2211-2212-2213-2214-2215-2216-2217-2218-2219-2220-2221-2222-2223-2224-2225-2226-2227-2228-2229-2230-2231-2232-2233-2234-2235-2236-2237-2238-2239-2240-2241-2242-2243-2244-2245-2246-2247-2248-2249-2250-2251-2252-2253-2254-2255-2256-2257-2258-2259-2260-2261-2262-2263-2264-2265-2266-2267-2268-2269-2270-2271-2272-2273-2274-2275-2276-2277-2278-2279-2280-2281-2282-2283-2284-2285-2286-2287-2288-2289-2290-2291-2292-2293-2294-2295-2296-2297-2298-2299-2300-2301-2302-2303-2304-2305-2306-2307-2308-2309-2310-2311-2312-2313-2314-2315-2316-2317-2318-2319-2320-2321-2322-2323-2324-2325-2326-2327-2328-2329-2330-2331-2332-2333-2334-2335-2336-2337-2338-2339-2340-2341-2342-2343-2344-2345-2346-2347-2348-2349-2350-2351-2352-2353-2354-2355-2356-2357-2358-2359-2360-2361-2362-2363-2364-2365-2366-2367-2368-2369-2370-2371-2372-2373-2374-2375-2376-2377-2378-2379-2380-2381-2382-2383-2384-2385-2386-2387-2388-2389-2390-2391-2392-2393-2394-2395-2396-2397-2398-2399-2400-2401-2402-2403-2404-2405-2406-2407-2408-2409-2410-2411-2412-2413-2414-2415-2416-2417-2418-2419-2420-2421-2422-2423-2424-2425-2426-2427-2428-2429-2430-2431-2432-2433-2434-2435-2436-2437-2438-2439-2440-2441-2442-2443-2444-2445-2446-2447-2448-2449-2450-2451-2452-2453-2454-2455-2456-2457-2458-2459-2460-2461-2462-2463-2464-2465-2466-2467-2468-2469-2470-2471-2472-2473-2474-2475-2476-2477-2478-2479-2480-2481-2482-2483-2484-2485-2486-2487-2488-2489-2490-2491-2492-2493-2494-2495-2496-2497-2498-2499-2500-2501-2502-2503-2504-2505-2506-2507-2508-2509-2510-2511-2512-2513-2514-2515-2516-2517-2518-2519-2520-2521-2522-2523



HCl	hydrochloric acid
HPLC-RT	HPLC retention time

THF	tetrahydrofuran
TFA	trifluoroacetic acid

# EXAMPLES:

[084] If not described otherwise, the examples mentioned here are racemates.

[085] The HPLC or LCMS conditions used for characterization were as follows: HPLC and HPLC-MSD systems from Agilent Technologies of the series 1100 with DAD detector, Merck Purospher column (3  $\mu$ , 2 x 55 mm), column temperature: 30°C, wavelength: 220 nm, flow: 0.5 ml/min, gradient: from 95% water (0.05% TFA)/5% acetonitrile in 4 min to 5% water (0.05% TFA)/95% acetonitrile, then the column was kept at 5% water (0.05% TFA)/95% acetonitrile for 1.5 min.


[086] The values marked by \* were determined under the following conditions: HPLC-MSD system from Agilent Technologies of the series 1100 with DAD detector, Nucleosil column (C-18, 5 $\mu$ , 4x125 mm), column temperature: 40°C, wavelength: 220 nm, flow: 0.65 ml/min, gradient: 95% water [water/acetonitrile 9:1 with 0.1% TFA]/5% acetonitrile [water/acetonitrile 1:9 with 0.1% TFA] for 2 min, then to 5% water/95% acetonitrile in 10 min, then the column was kept at 5% water/95% acetonitrile for 5 min.

[087] For the characterization of the final compounds, the HPLC retention time and the result of the mass-spectroscopic investigation, which was carried out separately, are given.

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

**[088] Example 1: (rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine hydrochloride**



**[089] a) exo/endo-3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-5-ylamine and exo/endo-(3a,4,5,6,7,7a-hexahydro-3H-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine**

10 g of exo-5-isothiocyanate-5,6-dihydroenedodicyclopentadiene (Maybridge international) were dissolved in 61 ml of formic acid and the solution was boiled under reflux for 45 hours. After cooling, a black precipitate was filtered off and the filtrate was concentrated. The residue was diluted with water, and 10 g of sodium hydroxide were slowly added in the presence of heat. The mixture was then cooled to room temperature, extracted three times with toluene. The combined organic phases were dried using  $MgSO_4$ , the  $MgSO_4$  was filtered off and the filtrate was concentrated. The residue was distilled and afforded 3.38 g of a clear oil.

HPLC-RT=3.15 min; MS (CI<sup>+</sup>): 150 (M+H)<sup>+</sup>

**[090] b) (exo/endo)-octahydro-4,7-methanoinden-5-ylamine**

3.3 g of the double-bond isomer mixture from 1a) were dissolved in 30 ml of methanol. 0.5 g of palladium on carbon (10%), as catalyst, was added thereto and the mixture was hydrogenated under a hydrogen atmosphere for 4 h. The catalyst was then filtered off, washed with methanol and the filtrate was concentrated. After drying under a high vacuum, 3 g of product were obtained as a clear oil.

HPLC-RT=3.33 min; MS (ES<sup>+</sup>): 152 (M+H)<sup>+</sup>


LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[091] c) exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine hydrochloride

A solution of 3 g of the exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine from 1 b) and 2.15 g of pyridin-3-carbaldehyde in 200 ml of toluene were heated to boiling for 5 hours in a water separator after addition of a catalytic amount of p-toluenesulfonic acid. After allowing to stand overnight at room temperature, the solvent was distilled off. The residue was dissolved in 150 ml of methanol and 0.91 g of sodium borohydride were then added in small portions to the ice-cooled solution with stirring. The mixture was stirred at room temperature for several hours and then rendered strongly acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was dissolved in water and rendered alkaline using potassium carbonate solution. The mixture was then extracted three times with EA, and the combined extracts were dried over magnesium sulfate, filtered, and adjusted to pH 1-2 using ethereal HCl. The solvent was then decanted off from the precipitated product and the residue was dissolved in ethanol in the warm. After cooling, the product was precipitated using ether. 2.85 g of pale crystals were obtained.  
HPLC-RT=3,15, MS (ES+) : 243.2 (M+H)<sup>+</sup>

[092] **Example 2: (+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine hydrochloride**



[093] a) (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine and (-)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine

The title compounds can be obtained enantiomerically pure starting from racemic (exo/endo)-octahydro-4,7-methanoinden-5-ylamine from Example 1 b) either by means

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

of chromatography on chiral columns or by crystallization using chiral acids. Using Z-valine, for example, (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine can be obtained as follows:

[094] a1) (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine by resolution using Z-valine

A mixture of 50 g of racemic (exo/endo)-octahydro-4,7-methanoinden-5-ylamine from Example 1b) and 83.2 g of Z-valine in 1.5 l of tetrahydrofuran was heated to boiling. The clear solution was allowed to cool to ambient temperature in the course of 3 hours. It was then stirred at ambient temperature for a further 20 hours. The precipitate was filtered off with suction, washed with 50 ml of tetrahydrofuran, and dried at 40°C in a drying oven. 55 g of the salt of 65% diastereomeric purity were obtained.

After recrystallizing three times from one liter of tetrahydrofuran in each case, 30 g of the salt of > 95% diastereomeric purity were obtained.

[095] 1 N sodium hydroxide solution was added to a suspension of 9.4 g of the above Z-valine salt in 30 ml of toluene and water until the pH remained constant at 11. In the course of this, the solid went into solution. The phases were separated and the water phase was extracted a further three times using 10 ml of toluene in each case. The combined toluene phases were dried over sodium sulfate, and the toluene was distilled off in vacuo.

[096] It was possible to react the amine thus obtained (3.3 g, enantiomeric purity > 95%) without further purification.


[097] a2) (-)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine by resolution using Z-D-valine:

The (-) enantiomer was obtained analogously to the above procedure using Z-D-valine. Thus, starting from 500 mg of (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine

A solution of 1.4 g of the (+)-exo/endo-configuration octahydro-4,7-methanoindene-5-ylamine from 2a1) and 1 g of pyridin-3-carbaldehyde in 100 ml of toluene was heated to boiling for 5 hours in a water separator after addition of a catalytic amount of p-toluenesulconic acid. After allowing the solution to stand overnight at room temperature the solvent was distilled off. The residue was dissolved in 75 ml of methanol and 0.42 g of sodium borohydride were then added in small portions to the ice-cooled solution. The mixture was stirred for several hours at room temperature, allowed to stand overnight and then rendered strongly acidic using excess methanolic HCl. The crystallization initiated by trituration was completed overnight in a refrigerator. After decanting off the solution, the residue was taken up with aqueous potassium carbonate solution and extracted three times with EA at pH 11, and the combined extracts were dried over magnesium sulfate, filtered, and adjusted to pH 1-2 using methanolic HCl. The product was then precipitated by addition of ether. 1.1 g of pale crystals were obtained.

[099] HPLC-RT=3.20; MS (CI<sup>+</sup>) : 243.3 (M+H)<sup>+</sup>; [α]<sub>Na</sub> 589 nm: +34,6° in ethanol

**[0100] Example 3: (-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethanamine hydrochloride**




[0101] A solution of 1.4 g of (-)-exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine from 2a2) and 1 g of pyridin-3-carbaldehyde in 100 ml of toluene were heated to boiling for 5 hours in a water separator after addition of a catalytic amount of p-toluenesulfonic acid. The solvent was distilled off after allowing the solution to stand overnight at room temperature. The residue was dissolved in 75 ml of methanol and 0.42 g of sodium borohydride was then added in small portions with stirring to the ice-cooled solution. The mixture was stirred for several hours at room temperature, allowed to stand overnight and was then rendered strongly acidic using excess methanolic HCl. The crystallization initiated by trituration was completed overnight in a refrigerator. After decanting off the solution, the residue was taken up with aqueous potassium carbonate solution and extracted three times with EA at pH 11, and the combined extracts were dried over magnesium sulfate, filtered and adjusted to pH 1-2 using methanolic HCl. The product was then precipitated by addition of ether. 1.1 g of pale crystals were obtained.

[0102] HPLC-RT=3.12; MS (Cl<sup>+</sup>) : 243.3 (M+H)<sup>+</sup>; [α]<sub>D</sub><sup>25</sup>Na 589 nm : -32.5° in ethanol

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

**[0103] Example 4: exo/endo-furan-2-ylmethyl-(octahydro-4,7-methanoinden-5-yl)-amine hydrochloride**



[0104] 200 mg of the exo/endo-configuration octahydro-4,7-methanoinden-5-ylamines from Example 1 b), 127 mg of 2-furaldehyde, and 101 mg of p-toluenesulfonic acid were dissolved in 20 ml of toluene (anhydrous) and the solution was boiled under reflux for 4 hours. After allowing the solution to stand overnight at room temperature, the solvent was distilled off. The residue was dissolved in 15 ml of methanol and 0.6 g of sodium borohydride were then added to the ice-cooled solution in small portions with stirring. The mixture was stirred for several hours at room temperature and then rendered acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was taken up in 2 N NaOH and extracted three times with EA. The combined organic phases were acidified using methanolic HCl and concentrated. The oily residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% TFA). The clean fractions were combined, the acetonitrile was removed in a rotary evaporator, and the mixture was adjusted to pH 11 using potassium carbonate and extracted with EA. The combined EA phases were dried using MgSO<sub>4</sub> and concentrated after filtering off the MgSO<sub>4</sub>. The residue was taken up with 2 N hydrochloric acid and freeze-dried. 119 mg of the hydrochloride were obtained as a white solid.


[0105] HPLC-RT=3.60 min; MS (ES<sup>+</sup>): 232.2 (M+H)<sup>+</sup>

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

**[0106] Example 5: exo/endo-(octahydro-4,7-methanoinden-5-yl)-thiophen-2-ylmethylamine hydrochloride**



[0107] 200 mg of exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine from Example 1b), 148 mg of thiophene-2-aldehyde and 101 mg of p-toluenesulfonic acid were dissolved in 20 ml of toluene (anhydrous) and the solution was boiled under reflux for 4 hours. After allowing the solution to stand overnight at room temperature, the solvent was distilled off. The residue was dissolved in 15 ml of methanol and 0.06 g of sodium borohydride were then added to the ice-cooled solution in small portions with stirring. The mixture was stirred for several hours at room temperature and was then rendered acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was taken up in 2 N NaOH and extracted three times with EA. The combined organic phases were acidified with methanolic HCl and concentrated. The oily residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% TFA). The clean fractions were combined, acetonitrile was removed in a rotary evaporator, and the mixture was adjusted to pH 11 using potassium carbonate and extracted with EA. The combined EA phases were dried using MgSO<sub>4</sub> and concentrated after filtering off the MgSO<sub>4</sub>. The residue was taken up with 2 N hydrochloric acid and freeze-dried. 61 mg of the hydrochloride were obtained as a white solid.


[0108] HPLC-RT=3.84 min; MS (CI<sup>+</sup>): 248.3 (M+H)<sup>+</sup>

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000



**[0109] Example 6: exo/exo-(octahydro-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine hydrochloride**



**[0110] a) Octahydro-4,7-methano-inden-5-ole**

25 g of tricyclo[5.2.1.0 (2,6)]decan-8-one (Aldrich) were dissolved in 100 ml of methanol and treated with 6.3 g of solid sodium borohydride at room temperature with slight cooling and stirring in portions in the course of 2 h. The mixture was then stirred for 2 h and allowed to stand overnight. About 40 ml of 2 N HCl were then added dropwise with cooling, followed by 20 ml of water. The mixture was concentrated, the residue was treated with ethyl acetate, and the ethyl acetate phase was washed once with water and once with sodium hydrogen carbonate solution. After drying using magnesium sulfate, the ethylacetate phase was filtered and concentrated. 26 g of oil remained, which was purified by vacuum distillation. 20.7 g of an oily liquid were obtained (bp<sub>0.5</sub> 76°C).

[0111] HPLC-RT=4.55 min; MS (CI<sup>+</sup>): 134.8 (M-OH)<sup>+</sup>

**[0112] b) 2-(Octahydro-4,7-methanoinden-5-yl)-isoindole-1,3-dione**

1.7 g of diethyl azodicarboxylate diluted with 5 ml of THF were added with stirring to a solution of 1.66 g of octahydro-4,7-methanoinden-5-ole from 6 a), 1.47 g of phthalimide and 2.62 g of triphenylphosphine in 15 ml of THF. After standing overnight, the reaction mixture was evaporated, the residue was stirred with ether, the precipitate was filtered off with suction, and the filtrate was concentrated. The residue was purified on silica gel/toluene. 1.36 g of a yellow oil were obtained.

[0113] HPLC-RT=5.82 min; MS (CI<sup>+</sup>): 282.2 (M+H)<sup>+</sup>

LAW OFFICES

[0114] c) exo/exo-octahydro-4,7-methanoinden-5-ylamine

0.4 g of hydrazine hydrate was added dropwise to a solution of 1.12 g of 2-(octahydro-4,7-methanoinden-5-yl)-isoindole-1,3-dione from 6 b) and 15 ml of ethanol and stirred at 65°C for 2 h. The mixture was then adjusted to pH 1-2 using conc. HCl and treated with 10 ml of ethanol. The precipitate was filtered off and the filtrate was concentrated. The residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% trifluoroacetic acid). After freeze-drying, 567 mg of product were obtained as trifluoroacetate. Treatment with sodium hydroxide solution and ethyl acetate yielded 322 mg of the free amine.

[0115] HPLC-RT=3.47 min; MS (CI+): 152.0 (M+H)<sup>+</sup>

[0116] d) exo/exo-(octahydro-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine hydrochloride


A solution of 332 mg of the exo/exo-configuration octahydro-4,7-methano-indenylamine from 6 c) and 215 mg of pyridin-3-carbaldehyde in 20 ml of toluene were heated to reflux for 5 hours after addition of a catalytic amount of p-toluenesulfonic acid. After allowing the solution to stand overnight at room temperature the solvent was distilled off. The residue was dissolved in 15 ml of methanol and 91 mg of sodium borohydride were then added in small portions to the cooled solution with stirring. The mixture was stirred at room temperature for several hours and then rendered strongly acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was taken up with 2 N sodium hydroxide solution. After extracting three times with EA, the combined extracts were concentrated, and the residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% TFA). The product-containing fractions were combined, freeze-dried, and again purified by HPLC. The clean fractions were combined, the acetonitrile was removed on a rotary evaporator, and the residue was adjusted to pH 11 using potassium carbonate and extracted with EA. The combined EA phases were dried using MgSO<sub>4</sub> and concentrated after filtering off the MgSO<sub>4</sub>. The residue was taken up with 2 N hydrochloric acid and freeze-dried. 35 mg of the hydrochloride were obtained as a white solid.

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[0117] HPLC-RT=3.25 min, MS (ES<sup>+</sup>): 243.1 (M+H)<sup>+</sup>

**[0118] Example 7: exo/endo-(3-chloro-5-trifluoromethylpyridin-2-ylmethyl)(octahydro-4,7-methanoinden-5-yl)-amine trifluoroacetic acid salt**




[0119] 0.5 mmol of exo/endo-octahydro-4,7-methanoindene-5-ylamine from Example 1 b), 0.5 mmol of 3-chloro-5-trifluoromethylpyridin-2-carbaldehyde, 140  $\mu$ l of triethylamine, and 10 ml of dichloromethane were introduced, 250  $\mu$ l of a 1-molar solution of titanium tetrachloride in toluene were added dropwise and the mixture was stirred at room temperature for 24 h. 1.5 ml of a 1-molar solution of sodium cyanoborohydride in THF were then slowly added and the mixture was stirred at room temperature for 30 min. The mixture was then treated with 15 ml of 2 N NaOH and stirred for 15 min. The solid was filtered off and washed with water. 30 ml of EA were added to the filtrate, the mixture was shaken and then the organic phase was separated off. After drying, the mixture was concentrated and the residue was purified by means of preparative HPLC (RP18, gradient acetonitrile/water 30%  $\rightarrow$  90%, with 0.1% TFA in both components). After freeze-drying, 4.7 mg were obtained as a white solid.

[0120] HPLC-RT= 11.23 min\*, MS (ES<sup>+</sup>): 345.2 (M+H)<sup>+</sup>

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

**[0121] Example 8: exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)-pyridin-3-ylmethyl-amine hydrochloride**



**[0122] a) bis-(3-chloro-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-yl)-diazene N,N'-dioxide**

3.34 g of isoamyl nitrite were added to a solution of 3.56 g of benzonorbornadiene [L. Friedman and F.M. Logullo, J. Org. Chem. 34: 3089-3092, (1969)] in 6 ml of glacial acetic acid and 6 ml of ethanol. 8.5 ml of a 15% strength solution of hydrogen chloride gas in ethanol were then added dropwise. The resulting suspension was stirred at room temperature for a further 2½ hours and then treated with 20 mg of diisopropyl ether. The solid was filtered off after further stirring for 30 minutes. A pale crystalline solid; m.p. 187 - 188°C, was obtained.

MS (FAB): 415.1 (M+H)<sup>+</sup>

**[0123] b) (exo)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-ylamine**

3 g of bis-(3-chloro-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-yl)-diazene N,N'-dioxide were suspended in 150 ml of methanol and hydrogenated in an autoclave with hydrogen at 100 bar and 100°C for 20 hours using Raney nickel catalyst. After filtering off the catalyst, the solvent was evaporated. The residue was treated with water, rendered strongly alkaline with NaOH, and extracted repeatedly with methyl tert-butyl ether. After drying the organic phases, the desired amine was obtained as a pale yellow liquid.

[0124] MS (ES<sup>+</sup>): 160.0 (M+H)<sup>+</sup>

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[0125] c) exo/endo-decahydro-1,4-methanonaphthalen-2-ylamine

A solution of 1 g of exo/endo-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-ylamine in 10 ml of methanol and 30 ml of 2 N hydrochloric acid was hydrogenated in an autoclave with hydrogen at 100 bar and 90°C for 10 hours using 0.4 g of RuO<sub>2</sub>. After separating off the catalyst, the solution was evaporated to half the original volume. The aqueous solution thus obtained was rendered strongly alkaline with 10 N NaOH and extracted repeatedly with methyl tert-butyl ether. After drying and evaporating the solvents, exo/endo-decahydro-1,4-methanonaphthalen-2-ylamine was obtained as a colorless oil, which was stored under argon.

[0126] MS (CI<sup>+</sup>): 166.2 (M+H)<sup>+</sup>

[0127] d) exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)-pyridin-3-ylmethylamine hydrochloride

A solution of 0.71 g of pyridin-3-aldehyde and 1.1 g of exo/endo-decahydro-1,4-methanonaphthalene-2-ylamine in 40 ml of toluene was boiled under reflux for 4 hours after addition of a small catalytic amount of p-toluenesulfonic acid (3-5 mg) and the solvent was then distilled off. After dissolving the oily residue in about 30 ml of anhydrous methanol, the solution was treated in portions with 0.335g of sodium borohydride with cooling and stirring, stirred at room temperature for a further 2 hours, and allowed to stand overnight. The solution was then rendered acidic using a solution of HCl in methanol, the precipitate was filtered off, and the solvent was evaporated. The residue was recrystallized from ethanol, m.p. 283-285°C.







[0128] HPLC-RT=3.38 min, MS (CI<sup>+</sup>): 257.4 (M+H)<sup>+</sup>

[0129] The compounds described below were prepared from carbonyl derivatives known from the literature and the appropriate amines, analogously to the example indicated:

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

Table 1

Example		Salt	Analogously to Example	MS	HPLC RT [min]
9		HCl	5	ES+ 232.2 (M+H)+	3.73
10		HCl	5	ES+ 260.2 (M+H)+	4.00
11		HCl	5	Cl+ 248.0 (M+H)+	3.88
12		HCl	5	ES+ 243.1 (M+H)+	3.63
13		HCl	5	Cl+ 243.0 (M+H)+	3.14
14		HCl	5	Cl+ 232.2 (M+H)+	3.05

202-408-4000

LAW OFFICES








FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

15		HCl	5	Cl+ 232.1 (M+H)+	3.05
16		HCl	5	ES+ 231.2 (M+H)+	3.76
17		HCl	5	ES+ 249.1 (M+H)+	3.47
18 (racemate)		HCl	5	Cl+ 244.1 (M+H)+	3.39
19 (+)		HCl	2	Cl+ 244.2 (M+H)+	3.41
20 (-)		HCl	3	Cl+ 244.2 (M+H)+	3.40
21		HCl	8	ES+ 258.1 (M+H)+	3.65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000




22		HCl	5	ES+ 244.1 (M+H)+	3.33
23		HCl	5	ES+ 244.2 (M+H)+	3.43
24		HCl	5	Cl+ 232.1 (M+H)+	3.42
25	 mixture 	HCl	5	Cl+ 241.2 (M+H)+	3.07
26		TFA	7	ES+ 330.1 (M+H)+	11.78*
27		TFA	7	ES+ 274.2 (M+H)+	9.75*

202-408-4000

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N.W.  
WASHINGTON, DC 20005  
202-408-4000



28		TFA	7	ES+ 293.1 (M+H)+	10.03*
29		TFA	7	ES+ 356.3 (M+H)+	11.18*
30		TFA	7	ES+ 319.3 (M+H)+	9.99*

## PHARMACOLOGICAL DATA

### Example 31: Description of the Caco 2 model

[0130] The Caco-2 cell line was acquired from the American Type Culture Collection (ATCC) and maintained in Dulbecco's Modified Eagle medium (high glucose content), supplemented with nonessential amino acids, L-glutamine, penicillin/streptomycin, and 10% strength fetal calf serum. The Caco-2 cell line was then kept in an incubator under a 10% strength CO<sub>2</sub> atmosphere at 95% strength relative humidity and 37°C. The cells were grown in cell culture flasks (175 cm<sup>2</sup>). For the transport investigations, the Caco-2 cells were inoculated into polycarbonate cell culture inserts (costar Transwells®, pore size: 3 µm, area: 4.71 cm<sup>2</sup>) at a cell density of 6.5 x 10<sup>4</sup> cells/cm<sup>2</sup> and incubated in six-well culture plates with a change of medium after 4 and 8 days and then every second day thereafter. 21 to 25 day-old monolayers were used for the experiments.

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[0131] In each test series, a 21 day-old monolayer was tested for its properties with <sup>3</sup>H-dextran as a permeability marker. The value of the transfer rate (cumulative) had to be in the range of 2% after 120 min.

[0132] After eliminating the growth medium from the apical and the basolateral side, the monolayers were rinsed with the transport buffer (Hank's balanced salt solution, pH 7.8; containing 2.8 g/l glucose). The cells were then equilibrated for 15 min at 37°C under a 10% strength CO<sub>2</sub> atmosphere. The HBSS buffer was then removed. The test compounds were dissolved in a mixture of HBSS buffer and DMSO and added to the apical buffer such that a 1% strength (v/v) DMSO solution resulted. The test concentration in the first experiment was 1 mM, and 100 µM in the second. The experiments were carried out at 37°C and started by the addition of 1.5 ml of test solution to the donor side (apical). Transport buffer without compound was added to the recipient side (basolateral, 2.5 ml). At various points in time, samples were taken from the basolateral side (1 ml) and replaced by fresh buffer solution at 37°C. Apical samples were taken at the start and at the end (120 min) in order to determine the recovery rate of the compounds by means of these concentrations and the cumulative basolateral concentration.

[0133] The compounds were analyzed by means of HPLC.

[0134] The apparent permeability coefficient ( $P_{app}$ ) was calculated by means of the following equation:

$$P_{app} = \frac{d_c \cdot V}{d_t \cdot A \cdot c_o}$$

[0135] wherein  $d_c/d_t$  is the flow through the monolayer (µg of compound/ml x s), V is the liquid volume in the collection chamber (ml), A is the surface area size of the monolayer (cm<sup>2</sup>) and  $c_o$  is the initial concentration (µg of compound/ml) in the donor chamber. The flow through the monolayer was calculated from the cumulative basolateral concentration at the appropriate point in time with the aid of the initially linear data curve (linear up to 60 min). The respective determinations were made in





LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[illegible]

Results:

[0136] Table 2

		Absorbed fraction [%]
Example 1		100
Example 18		100
S 3226		<5
S 2120		<1

2025 RELEASE UNDER E.O. 14176

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

[0137] Compared with the NHE3 active compounds of the acylguanidine type known from the literature (J.-R. Schwark et al. Eur. J. Physiol., 436:797 (1998)), the compounds of the formula (I) or (I a) show a clearly superior ability to cross the membrane.

### Example 32: Description of the NHE activity measurements

[0138] Most of the molecular biology techniques follow protocols from the works "Current Protocols in Molecular Biology" (Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A. and Struhl, K., eds., John Wiley & Sons) and "Molecular Cloning: A Laboratory Manual" (Sambrock, J., Fritsch, E.F., and Maniatis, T., Cold Spring Harbor Laboratory Press (1989)).

[0139] In the context of our studies, stably transfected cell lines were produced that in each case expressed one of the following NHE subtypes: Human NHE1 (Sardet *et al.* Cell 56:271-280 (1989)), rabbit NHE2 (Tse *et al.* J. Biol. Chem. 268:11917-11924(1993)), human NHE3 (Brant *et al.* Am. J. Physiol. 269 (Cell Physiol. 38) C198-C206 (1995)), or rat NHE3 (Orlowski *et al.*; J. Biol. Chem. 267:9331-9339 (1992)).

[0140] The cDNA clones of the respective NHE subtypes obtained by Prof. Pouyssegur were, after addition of suitable linker sequences, cloned into the expression plasmid pMAMneo (obtainable, for example, via CLONTECH, Heidelberg) such that the recognition sequence for the restriction endonuclease NheI of the plasmid was approximately 20 – 100 Basepairs before the start codon of the respective NHE subtype and that the total coding sequence was present in the construct. In the case of the human NHE3 obtained from human kidney mRNA by means of RT-PCR, the RT-PCR primers were selected such that the cDNA band obtained had cleavage sites suitable for pMAMneo at its ends.

[0141] Using the "calcium phosphate method" (described in chapter 9.1 of "Current Protocols in Molecular Biology"), the NHE-deficient cell line LAP1 (Franchi *et*

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

*al.*; Proc. Natl. Acad. Sci. USA 83:9388-9392 (1986)) was transfected with the plasmids that received the respective coding sequences of the NHE subtypes. After selection for transfected cells by means of growth in G418-containing medium (only cells which have obtained a neogene by transfection can survive under these conditions), selection was made for functional NHE expression. For this, the "Acid Load" technique described by Sardet was used (Sardet *et al.*; Cell 56:271-280 (1989)). Unlike untransfected LAP1 cells, cells that express a functional NHE subtype can also compensate for the acidification carried out in this test in the absence of  $\text{CO}_2$  and  $\text{HCO}_3^-$ . After repetition of the "Acid Load" selection several times, the surviving cells were inoculated into microtiter plates such that one cell per well should occur statistically. Under the microscope, after approximately 10 days, a check was made to estimate how many colonies were growing per well. Cell populations from individual colonies were then investigated using the XTT proliferation kit (Boehringer Mannheim) with respect to their survival ability after "Acid Load". The best cell lines were used for further tests and, to avoid a loss of the transfected sequence, cultured in G418-containing medium under continuous selection pressure.

[0142] To determine  $\text{IC}_{50}$  values for the inhibition of the individual NHE subtypes by specific substances, a test developed by S. Faber (Faber *et al.*; Cell. Physiol. Biochem. 6:39-49 (1996)), which is based on the "Acid Load" technique, was slightly modified.

[0143] In this test, the recovery of the intracellular pH ( $\text{pH}_i$ ) after an acidification, which in the case of functional NHE occurs even under bicarbonate-free conditions, was determined. For this, the  $\text{pH}_i$  was determined using the pH-sensitive fluorescent dye BCECF (Calbiochem, where the precursor BCECF-AM is employed). The cells were first loaded with BCECF. The BCECF fluorescence was determined in a "Ratio Fluorescence Spectrometer" (Photon Technology International, South Brunswick, N.J., USA) at excitation wavelengths of 505 and 440 nm and an emission wavelength of

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

535 nm and converted into the  $pH_i$  by means of calibration curves. Differing from the described protocol, the cells were incubated in  $NH_4Cl$  buffer (pH 7.4) even in the case of the BCECF loading ( $NH_4Cl$  buffer: 115 mM NaCl, 20 mM  $NH_4Cl$ , 5 mM KCl, 1 mM  $CaCl_2$ , 1 mM  $MgSO_4$ , 20 mM Hepes, 5 mM glucose, 1 mg/ml BSA; a pH of 7.4 is established using 1 M NaOH). The intracellular acidification was induced by addition of 975  $\mu$ l of an  $NH_4Cl$ -free buffer to 25  $\mu$ l aliquots of the cells incubated in  $NH_4Cl$  buffer. The rate of pH recovery that followed acidification was recorded for 2 minutes in the case of NHE1, for 5 minutes in the case of NHE2, and for 3 minutes in the case of NHE3. For the calculation of the inhibitory potency of the tested substances, cells were first investigated in buffers in which a complete pH recovery occurred and in buffers in which no pH recovery at all took place. For the complete pH recovery (100%), the cells were incubated in  $Na^+$ -containing buffer (133.8 mM NaCl, 4.7 mM KCl, 1.25 mM  $CaCl_2$ , 1.25 mM  $MgCl_2$ , 0.97 mM  $Na_2HPO_4$ , 0.23 mM  $NaH_2PO_4$ , 5 mM hepes, 5 mM glucose, a pH of 7.0 is established using 1 M NaOH). For the determination of the 0% value, the cells were incubated in an  $Na^+$ -free buffer (133.8 mM choline chloride, 4.7 mM KCl, 1.25 mM  $CaCl_2$ , 1.25 mM  $MgCl_2$ , 0.97 mM  $K_2HPO_4$ , 0.23 mM  $KH_2PO_4$ , 5 mM hepes, 5 mM glucose, a pH of 7.0 is established using 1 M NaOH). The substances to be tested were prepared in the  $Na^+$ -containing buffer. The recovery of the intracellular pH at any tested concentration of a substance was expressed as a percent of the maximum recovery. From the percentage values of the pH recovery, the  $IC_{50}$  value of the respective substance was calculated by means of a Sigma Plot program for the individual NHE subtypes.

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000

### NHE activity

### Table 3

Example	Rat NHE3 IC <sub>50</sub> [μM]
1	0.34
5	2.9
6	2.1

[illegible]

## LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, DC 20005  
202-408-4000